

707a
707d

FROM
THE LIBRARY
OF
SIR WILLIAM OSLER, BART.
OXFORD

7666 k

January 13th

P. M. on a case of Phthisis. (See W. Jenner)
Much emaciated. Attention drawn
to the shape of the thorax, which was un-
usually long, flattened, the lower ribs
extending to within a short distance
of the costal dim. Sir W. remarked
that this was a form of thorax often
seen with congenitally small lungs,
and nature took this method of
adapting the parietes to them.
The upper ribs were oblique & the
intercostal spaces wide, but from
the fourth they began to slope extre-
mely, till about the 8 or 9th they
became almost vertical: the cost-
al cartilages as they ascended to their
insertion forming almost an acute
angle with. On opening the abdomen
the situation of the viscera was re-
marked upon: the liver being ~~slan~~ pla-
ced almost vertically and extending
in below the margin of the ribs. On each

side, when it came in contact with the costal cartilages, well marked patches of fat were seen. The Diaphragm was much arched the anterior portion being almost vertical & lying in close contact with the cartilages & the sternum. Attention was then drawn to the post mortem staining of the various parts and to the transudation of serum from the heart into the pericardium & to a slight degree into the pleura from the lungs. Open the heart always in the course of the circulation.

The de-colourized portions of the blood clot in the heart often afford important evidence in medico-legal inquiries with reference to either the original position of a body or as to whether it had been turned; as the colourless portions are always uppermost. Heart was flabby but otherwise healthy. Extensive pleural adhesions

all over both lungs. Right one pre-
sented at the apex signs of old standing
pneumonic affection(?), being indurated
in spots, channelled, small cavities
& studded here & there with caseous
masses. Throughout the rest of the
lung miliary tubercle was scattered
densely and Dr. W. remarked that
this was one of a very few cases he had
observed in which there was an
extensive deposit of tubercle, without
any great elevation of temperature
being, never above 100°. Some oedema
& emphysema were also observed
The left lung was small, shrunken
firm to the touch, though not curless,
looking very much like a lung comp-
ressed by fluid. The posterior portion
was channelled by an irregular cavity
of recent formation devoid of a lining
membrane strewed with numerous
bands & trabeculae

January 14th

P. M. on case of ^{(male) at 40} supposed lymph-
atic Leucemia. (Wilson 77)

Much emaciated & externally, large
glandular swelling, could be seen
in the right cervical and axillary
regions. The condition of the lymphatic
glands throughout the body was as
follows: Axillary & cervical of right side
formed a continuous chain, the
former being much matted together
making a lump about the size of a
hen's egg; the latter more isolated
None of any size in the corresponding
situations on the opposite side, nor
were there any in either groin
The sub-bronchial glands were
much enlarged, especially those
over the left bifurcation of the trachea,
which though completely pervious
yet had its caliber diminished
The lumen of the aorta was con-
siderably narrowed, hardly admitting

the little finger. The whole chain of retroperitoneal glands were enlarged, indurated & matted together and also formed adhesions with the caecum & pancreas

In section all these glands presented with slight modifications, the same appearance, they were firm to the touch, yet broke under the nail, in colour varying from a dirty white to a distinct yellow; traces of glandular structure sometimes visible capsule generally well marked & adherent. The variations were as follows: ^{A few of the smaller glands had a peculiarly tripe-like} One or two of the smallest ^{except with little or no cysts or a brown-mud kidney} and in the axilla had suppurated others could be easily enucleated. In the bronchial glands much pigment was present & in some a soft nucleus was observed. The retro-peritoneal glands differed only in one particular viz, that in one or two a peripheral line of suppurated

taken place between the band and
gland & the capsule

Lungs & diaphragms collapsed in position.
Apices of both the seat of tubercular deposit
(Fox) and tuberculous masses
were recalled throughout the organ.
Pigmentary connective tissue & inter-
lobular tissue between the nodules at the apices, in
which some central softening was
observed. Heart normal.

Spleen 3 (not very large) firm, albumi-
noid; sugar grains well marked
A patch triangular in form, white-
colour, and probably an infarct & site
at one extremity. Liver not enlarged
& apparently healthy.

Intestines. One typical tuberculous
ulcer & ulcers about one foot above the
ileo-caecal valve extending completely
round the gut ~~and~~ constricting its
diameter considerably. On the serous
surface of it some grey granulation
(tubercles) were found. Ulceration

not however for tuberculus charac-
was also found in the caecum, & much
a granulations on the serous coat ex-
ternally. Kidneys enlarged &
slightly albuminous

Dr Fox in remarking in this case
said, that, enlarged lymphatic
gland might arise from many
causes among which the chief
were; irritation, simple & specific,
scrophulous, hereditary, carcinoma
& the disease known as Lymph-
adenoma, This case was believed
to be one of the last named. There had
been no increase of white or corpus-
cles, but numerous small red ones
were in great excess. He had had albumi-
nuria and diarrhoea, but the chest-
disease, though diagnosed, was not
thought to be so extensive as I thought
it. Altogether Dr Fox inclined to the
belief that this was not a true case
of " Hodgkin disease " but a case of

general tubercular affection of the
lymphatics & to his mind the
similarity of the ^{appearance} affection in the
lungs, especially at the apices
favoured this view. The non-ex-
istence of adenoid growths in
other organs was also a strong ar-
gument. The Brain was healthy
and unaffected.

January 16

Out pat. B. Hill

Syphilitic orchitis

Testis uniformly swollen, epididym
not distinguishable. devoid of any
special secretions, differing in
this from the tuberculous and adu-
erous forms. May subside & then come
on again. Its final termination under
cure is ^{marked in} shrinking of the testis & imbe-
cility. Blend P. S. L. S.

Case of acute Eczema following a
scrape on the skin. In H. referred to
the emulsion of cod liver oil.

under which Eczema most frequently
arise among them the chief is the Gouty
In states of the system, dirt, cold, prostrated
power and the working in irritating
material is an active cause,

As regards the treatment. Wash well
first with soft water or water rendered
soft by boiling, then apply a powder of
3 parts Starch to one of Zinc, wrap the
limb in cotton wool and lay on
absolute rest

January 17

Dr. W. Jenner

In applying a hot fomentation, draw the patients clothes well up
about his anppits, place a thick fold
of blanket under the nates, then apply
your fomentation & cover with a piece
of old silk larger than the flannel
beneath, over this place an oiled fold
of blanket & over all fold the ends
of the blanket beneath his nates
This will keep warm for some hours

In taking observations with the
Aetherscope, in order that they
may be reliable you must repeat
them several times at the same
places with an interval of some
days between

In some women & in nearly all
children the voices do not
produce vibration enough to enable
you to feel the vocal primities. As
this is a very important aid in
the diagnosis of pneumonia from
pleurisy with effusion, what examina-
tion can you resort to with these classes
of patients? Dr. W. says we find it
in the displacement of organs. In
pneumonia the lung retains its own
bulk, but in a chest the sub-phrenic
gives a uniformly dull note behind &
the incise to the nipple in front. This
& a certain will produce some displa-
cement - should always be noted for

Dr. Keizer

Care of Syphilitic eruption
about the face & neck in a child 12
months old, suffering also from Thrush
and Diarrhea. He ordered $\text{gr} \frac{ss}{ii}$ of
Hydrog. c. exta. three times a day
and said that ⁱⁿ children diarrhea
was often stopped readily in the way
though at first it might aggravate
it a little. If the one dose was given
at night it would cause much
sleeping & diarrhea but - she could not
explain it if it were given three in the
day no ill effects follow &
Tinct. Copiva tried in Dropsy
Glycerine of Taraxacum in Eczema behind
the ears in children

Dr. Keizer

in referring to intestinal
obstructions said, "It was as culpable
to give cathartics on the one as in strangulated
hernia give opium which retards
the excellent spasm of the gut."

varm enematæ &c.

January 18th

Beneficial for Women. Colotomy. Mr Heath
This was a case of cancer in the rectum
not however producing complete obstruc-
tion, but as it was progressive and
the woman at present in good health
It was deemed advisable to operate
now, as by this means her chances
would be greater. Mr Heath made
a horizontal incision about 7 inches
in length the center of which cut a
vertical line. A saw was to pass
through the perineum third of the circle
dia. The various fasciæ having been
cut on a director, the bowel was exposed
As no feces were contained in it and
as they had been unable to pass up
up & dilated the bowel, some difficulty
was experienced in drawing it forward
Several folds of fat also impeded the
free access to the gut. Finally it was ob-
tained & before incising it two intestines

P. M. Scarlatinal droopy. Death from
dyspnea. Boy at 10

Anasarca general. 3 x x of milky
fluid in right pleural cavity. 3 iv
of mucus in the pericardium. Exam-
ination of this showed numerous
lymph corpuscles with. To inspect
all nucleated some multinnucle-
a few larger more granular cells
were also found

Right lung somewhat adenomatous,
pale, tough to the feel. Bronchi (in lobe
seemed thickened. Left lung.

On the surface of this was situated
- anteriorly & laterally - a large flattened
mass of fibrinous effusion, from $\frac{1}{4}$ to
 $\frac{1}{2}$ an inch in thickness, the exterior
being very firm, the interior more gel-
atinous & apparently more recent
organization. Evidences of tight compression

were visible beneath this, but otherwise the lungs did not present any marked features.

Heart normal. Large de-coloured clot in right auricle, fluid blood in right ventricle.

Liver healthy. Spleen soft & pale. Kidneys large. On section cortex and medulla of pale yellow colour. Medullary portion congested. Capsule easily detached. Surface of cortex pallid with numerous stellate veins scattered over it. A few pits & depressions were seen, perhaps the beginnings of pyramidal changes. The kidneys were good specimens for a stage not often seen ^(P.M.) after acute nephritis; the result of the attack was part of the organs apparently in such a condition that further detumescence changes or a complete recovery would depend on the state of bodily health.

Brain, pale, anemic. Surfaces on section "wet". Dura mater adherent.

January 21st

R. L. Ophthalmic in the morning
"Larson" case of indolent for dense
pupil "Scurvy of eye". Injury 7
years previously, by the explosion of
gunpowder, subsequent iritis, trau-
matic cataract, ~~injury~~ - Pain
& danger of sympathetic Ophthalmia
in the other eye, which had just re-
ceived an abrasion. Lodging in
the sclerotic beneath the tendon
of the internal Rectus a piece of
glass ~~had~~ about that eye was found
"Couper"

Congenital Cataract ^{.37 mm}
attempted solution, but found the
lens so hard & firm that extraction
was performed. Present operation to
tear with needles some remnants of
capsule left in the right eye

Care of Strabismus, internal,
Old man, gamekeeper, & troubled with
double vision;

U. C. H. "Finger"

January 30th

(Dubuque Fox)

Case of Apoplexy

Man at 75. well made, well nourished

History

Post mortem

Dura mater adherent
to the calvarium, which was unusually
dense and heavy. Sinuses normal
Surface of brain intensely hyperemic
and few spots of sub-arachnoid haem-
orrhage were found; one over the left
^{anterior lobe} temple, which corresponded ~~to~~ a
bruise received by the fall in the seizure
the second on the surface of the right
middle lobe, the third on the inner

surface of the left posterior lobe: the
remaining two occupied a position at
the under side of the cerebellar lobes.
The amount effused was not very
large, but in each case it extended
over a considerable area - in section
it was found deep in the sulci.
On slicing the hemispheres, the cen-
trum is marked externally, was
not here evident, though the vas-
cular prints were well defined.
On opening the right lateral ventricle
an extensive hemorrhage presented
itself, which on washing away the
clots was seen to have proceeded
from the body & hinder part of the
capsula striata and the anterior
segment of the Thalamus opticus, both
of which organs, together with the ad-
jacent structures, were much lacerated
and deranged. The septum was
also ruptured and clots with fluid
blood were found in the left ventricle.

le, the walls & floor of which were how-
ever intact. The haemorrhage had
also penetrated into the third ven-
tricle

The arteries were irregularly atherom-
alous, firm & fibrous rather than
calcareous

Nothing noticeable in other portions
of the organ

Heart, large, with a good deal of per-
icardial fat lying upon it

Right chambers, normal, with the
exception of slight opacity of the lining
membrane of the auricle


Left Auricle healthy. Ventricle much
hypertrophied, but the muscular fibres
apparently perfect, no traces of "aged
leaf patches"; indicated fatty changes
Mitral valves, somewhat thickened
and the orifice slightly narrowed
One or two nodular masses of yellowish
atheroma were attached near
the margin of one segment. Aortic


valves thickened slightly, seen one
commencing calcification


Lungs extensively collapsed and
adematous and at the base small
tracts of Pneumonia could be seen
distinct from the collapsed and
adematous lung by its want of tran-
slucency & slightly granular aspect.
Liver & Spleen firm but otherwise
normal.

Kidneys, small, markedly fibroid
Capsule detached with great difficulty,
inner surface granular & pale
It seems as if in these cases, the puer-
ery of the cortex, take place at the
entrance of a blood vessel only, the
round smooth granules, being
devoid of vessels. On section, cortex
reduced in up. pyramid within $\frac{1}{8}$
of an inch of the surface. Malpighi
bodies indistinct. No pyramidal
congestion, cystic degeneration in
spots

winded park is one of the few things here worthy of
unqualified praise & indeed is the only place
where we can take a walk with any comfort
the general surroundings of the city are so flat &
uninteresting - we can be well imagined







No 1 (continued)

especially in left lung the surface was
of a uniform colour and without these
cancerous masses. Portions of collapsed lung
in lower lobes, and there is a good deal of
blood and serum in dependent parts.

The bronchi contain an exceedingly tenacious mucus, and
some of the smaller tubes have the cancerous matter in
them. Spleen looks natural, pulp not increased
consistence good, Mal. corp. abundant.

Kidneys, look a little opaque, otherwise normal.
Liver looks normal. Stomach contains a quantity
of half digested food. The Cardiac end contains
much blood, the M. M. lining of a uniform red colour,
feels a little soft. Throughout the small intestine
the M. M. presents a patchy redness due to capillary
injection of the mucosa. In many places it is uniform.
Peyer's glands are ^{not} affected, neither are the cecum or
scrother. Large intestine, mucous membrane presents
the same appearance, no ulceration or involvement
of the glandular apparatus, merely congestion of the
mucosa. The Meckel's gland is a little enlarged
not haemorrhagic.

No 1

Male pig 6-7 months old. Numerous bloody
ecchymoses upon skin of back flanks and ears.
No scab, vesicular, pustular or pustular eruption
on open abdomen muscles of a pale colour, no point
of anal effusion. Intestines look tolerably normal
for serous surface. In chest, left lung is rather
inflated, ^{a little} fluid out by infl. about 3. Pleural
fluid in pericard. and there are a few flakes
of lymph in it. The surfaces of the membrane
a little cloudy. Heart bright chambers are distended
with blood, valves healthy, left a small clot in left
vent. walls & valves normal.

Right lung. almost the whole of the upper lobe, and
the ant. end of the lower is in a condition of granu-
lar pneumonia, redundant & airless, the pleura covering it being
irregular & rough. here & there ecchymotic. On section
it appears as if small cavernous spaces were scattered
through the tissue, they have rounded outlines ranging
from quite short to peas, opaque white in colour, and
in pressure may often be seen to be in connection with bronchi
and probably they can be squeezed as casts. The tissue
then is dark red; hepatized, in some areas

No 2,

male fig. 6-7 mounted. Skin of back & abdomen also about ears eccyrotic, in irregular patches, the other cutaneous affection. Pig. not disappearing. On sp abd, no fluid in cav. Sides look natural. In thorax, no fluid in pleura. Pericard contains a few diatoms and a thin layer of lymph gives the base, lighter, remaining in the visceral. The menbr are somewhat injected. Heart contains blood, right chamber small, cavity in left cut. Valves & valves normal. Lungs, extremely collapsed, very dark in colour and the lower lobes almost black. The upper part of ant lobe alone of normal colour. The bronchial tubes contain a large quantity of mucus, and are in tubes of 2nd char. than whole tissue, mucus noticed in the mucus. On skull, of the trachea a similar greyish white mucus lines the entire surface from into bronchi, he stripped in a part of the tube, it is then about 1/4" in thickness, especially, can't learn a much injected surface Bergall which looks a little roughened. This extends from the tubes of 2nd char. through the bronchi, lines the larynx & is a thin cloud, almost dark grey membrane covers the epiglottis the pillars of the fauces, the tonsils and extends into the posterior nares, coating the entire fund. and the septum. The m m beneath these latter parts is very much congested and of a dark purple colour, the mucus glands of the tongue are enlarged. Spleen looks natural, dark reddish brown, mucus, pulp not increased, connect-tissue. Liver looks very healthy, looks in form mid. full blood. full. Kidneys a little cloudy, but very natural looking. Stomach contains half digested food, non a little reddened, and with special to be seen. Small bowel contains a dirty fluid, feces. M. m. not congested. Peyer's patches, not much. A few small raised nodules congested spots seen a short dist from venter looking like inflamed spots glands beginning to ulcer on lips. In the caecum & colon are numerous small clashing spots ranging in size from 0 to 1/2. The smaller ones present a central white spot surrounded by a green or whitish hyperaemia. The larger ones have a dirty greyish yellow central though beginning to separate in some. also curr. 4/3 size of hypera. a few are seen still darker dot where a small white slightly thickened mass - a gland(?) is seen one of 1/2 a ring of congestion. The end of both tubes at the caecum & around colon. The m m below the spots present irregular areas of hyperaemia but looks more normal. The menbr of Semivale swollen slightly, thickened but not much of it all congested.

No 3.

Female by 4 months.

Large Intestine. The head of the ~~caecum~~ caecum looks dark from outside & is covered with E.C.C. the glands in its neigh- are of a dark purp. colour somewhat swollen & on section the cool part is the seat of retroa - the mucous membrane is entirely congested somewhat thickened & at one part covered with a uniform dirty membrane the rest of the caecum presents a large number of swollen glands those near the head are close together, may have merged together & present a dark discolourization on the

surface higher up -
the gut they are more
isolated & present brownish
yellow tops immediately
above the valve there are
two or 3 flat ulcers size
5 cent by 18" from
this part the wall is
thickened & presents
numberless confluent ^{after} ^{slough}
sloughs which are still
closely adherent to the
tissue beneath some
have separated leaving
flat ulcers. for the
next ft the intestine is
much congested covered
here & there with a
dirty grey membrane
the solitary glands are
swollen & here & there
is a distinct ulcer the
rest of the large intestine

is somewhat congested
but is not ulcerated
Lungs with except of
some collapse & one
small patch of inflammation
appear healthy. Beneath
the peric over the right
oricle is a uniform
layer of blood & portion
of the wall of this
cavity appear infiltrated
with blood the same
condition though not
so marked in the left
liver an intense
dark slate color which
is uniform throughout
Spleen not enlarged
consistence good
pulp moderate
seen in the sub
mucous tissue of the
rectum in a large

portions of its extent are
numerous hemorrhages
the intestine looking
quite black

Additional on No 3 Cap caecum presents a uniform grey tenace
when adhe mount over an ar. $\frac{1}{2}$ to $\frac{1}{4}$ " & beneath it the mucus is note
conspicuous. A few an. outlets mount & sculler over in the lower half with
round clear, whitish bodies. Flat dark on a scale in the center, a pale
sandy border which fades directly over into the darkness of m m.

The lips of some are depressed, & they lie in a shallow pit sur-
rounded by a pinkish red ring. 3-8-10 of these are often
mingled together. In the upper half of caecum they are small, the lips are
not so flat - and the little central opening is began in many to
expand. A few spots are seen like swollen set glands, though
but they are small. 7 or quite 2 feet above the valve the m m present
a rough leather like appearance the whole wall is stiff and firm, it
looks as if the m m were up to with a very slight elevation, which here
is there in beginning to rise & lay on irregularly, of dark & white
surface beneath. Scattered over this area are old numerous
patches as in caecum. Farther on in the gut 3rd & 4th foot the
mucus is dotted over with small spots to the m m. In many
of which the length is separating. The small collection mud, pellets
The terminal portion of the stem and the valves are not covered in flum
a grey patch extends to the tubercle is perfect, normal.

Case 4 Pig 4 months old. Had been ill

was very weak sluggish. Bod. temp. much increased
careful inspection fails to discover any rubeloid
hyper. rash. There are a few blotchy ecchymoses
and the ears present a purplish color. The animal was
bled to death. ^{slightly} Blood somewhat dark but looked natural.
Nothing abnormal in abdominal cavity. In thorax lungs
collapsed. Pericard. normal. Heart. healthy. Small partly
decalcified clot in left vent. Clotted, adherent to valve.

Lungs quite healthy a few patches of collapse noticed.
Trachea. Pharynx & tubes healthy. Spleen a little
larger than the others. but firm. pulp not increased
are wedge shaped haemorrhagic infarct on free edge
kidneys look healthy. Liver almost good. Lobules a little pale in
center. veins and gen appear normal. Stomach
contains recently ingested food. In. M. perfectly natural.
Nothing abnormal in small intest. Peyer's glands abundant,
and very long. not at all swollen or affected.

In the colon are numerous polyps, of a very remarkable appearance
at the ileo-caecal valve are four flat ~~spots~~ on the mesent. They have
merged in a line and extend 3" in the length of the gut. They
have a rounded apical border, a slightly irregular surface, which
is in 2 of the shaded black they present a concentric arrangement
of lines caused by small pieces of mucus around a central portion
one or two small masses in the caecum. Higher in the
colon are others, scattered, 6-8, the size of pennies perfectly
round, & present the peculiar concentric arrangement.
The central part is a little darker than the av. They
look like nuptial cups flattened out

Histological Examination


Liver ~~case 1~~ ^{In one case} ~~Dark slate colour~~. Traced portions show the cells stuffed with small fine granules, some of which look black, the general appearance is that of cells in condition of cloudy swelling. The nuclei are distinct & not obscured. F.M. in the field are innumerable granules and also cells, resembling in all particulars nuclei of hepat. corpuscles. In certain spots traced into the liver cells instead of being distinct and forming a shaggy column, appear as if merged together into a homogeneous granular mass in which the nuclei are irregularly scattered.

^{mesothelium} lymphatic gland (case 2). Portions after one, which appears of a uniform, ~~dark~~ purple colour, show an enormous ^{number} of blood corpuscles within the gland tissue; the lymph corpuscles are normal looking. A few giant cells are seen. At one spot near the center a collection of somewhat large corpuscles the true veg. granular int. corp. was seen surrounded by an irregular stratum.

Uterus. The peculiar carcinosomatous material which follows the branched tubes in the hepatic portion of my case 7 consists chiefly of small round translucent corpuscles, which, looking like free nuclei and about $1/3 - 1/2$ the size of colourless blood

corpuscles. They ^{tend to} adhere in clumps in stained preparation. &
occasionally 2-4 may be seen appressed in a slightly granular
matrix, a few were noticed to have a ring cell substance
about them. These elements comprise the bulk of the material
col in the tubes. Scattered here & there are larger corpuscles,
like dendrocytes and cells of per. epithelium. 1

In the majority of the cases the ^{masses} lymph glands
were swollen enlarged & tumefied, often of a
deep purple color, depending upon the exhaustion
and effluence into the cortex.



Experiments

1st Sept.

Saturday 7.30 P.M.

I a three month old ^{white} sow pig was
inoculated with lymph and blood obtained by
squeezing the portion of the skin - where it was ecchy-
motic - of Carl I and collecting the exudate on ivory points

The virus was performed in 2 places by scratching
the skin - on abd. and near groin - with the ivory point.
2d. appears all right. T 104.2. 3d. no change. cuts well looks
fine. 6th not so well. does not come to the side of the pen. & is
with difficulty stirred up. The huddle together. 7th a.m. T 105.4. Has
skin well, no diarrhoea. Plane of nerve has dried up. A diffuse cut-
aneous redness over the skin of belly, which disappears on pressure. a few
spots instead of circumscribed papules, irregular in outline. chief about
hair follicles. they are then more swollen & somewhat papular. They exist
in flanks and on belly. 8th. - - - - -

9th a.m. T 106.4. Skin on abd not so marked, but one or two faint
spots? visible. no effect of carb. niger. etc. they feed well

11th. appear very well cut in several. One or two
irregular papules, about no ecchy or blotches. or hyperaemia
T 107.

14th. T. 106.2. no change to note. animal feeds well.

17th. M. T 105.4. Animal looks well. Back somewhat
darker up. Skin looks harsh & dry. Skin of abdomen of a
dusky red colour, and covered with small slightly
elevated papules. Skin of back of ears also very dark
& in part ecchy-motic.

18th. T 102. Skin for whole body of a deep dusky
red colour, somewhat brighter on abdomen. It dis-
appears in most places on pressure, returning very slowly

when the pressure is removed. Over sternum & in
axilla the skin is in place ecchymotic. Backs of ears
very dark. Inguinal glands enlarged. animal
does not feed well & looks ill. Eyes bulging.

D, T 102, not so red, but skin b. No diarrhea. M M fed red

19th M. T 104° does not feed well no diarrhea E. T 105°

20th M. T 105° Not so red. Skin very hard. Lymph

glands most swollen. Examined the blood. no m
red corp with normal aggr. in clumps not in def solution. The color is off
somewhat increased & very little in eye. Schuller granules
on reds large & numerous, the white elements of them very distinct
a few fine parasites. No bacteria. Fibrin fib. not abund. P M T 105°

21st M. T 104° Redness much less, a few blotches on abdomen, but hard
disappears on pressure. P M 10 3/4

22nd 104.4 Rash little marked. Glands still swollen

23rd 104°

24th 104° 25th 103° 26th 106° No rash visible

27th 104.4 28th 104

Experiment II

Sept 12th 7.20 P.M.

Heck's white
a three months old serpig

was unincubated, by scratching & rubbing in the matter, with material obtained from one of the plagues of case IV. and punctured place on under surface of belly, with material for inoculation from intestine of case III

Pulse unaltered after operation 100. T 106.4 (This with

two separate thermometer). Nothing unusual noticed at the moment even until the evening of the 6th when the animal appeared so lively & lay towards the stern. 7th at the spot underneath of belly where above there is an elevated swelling underneath flattened on the top, rest of all unaltered. about 2" from it towards the right groin is an oval swelling, the size of a small plum, looking a little bluish, & feels like a swollen lymph gland. The other spot on the groin presents itself, any signs of reaction. A faint, just perceptible redness - noticeable in abdomen, and on inner surface of arms. a few small papules at cost of L. vis 8th. arm 9th. T 105.5. where no cuticleless nodules. Redness had disappeared. Papules not so marked. Feeds well. 5

11th Sides more marked by a flat tubercle swelling one or two papules on flank. Is hyperaemia a scabium appears quite well T 106°

14th T 107°. Little change. Site of puncture marked by elevated spot, and a lymph gland in neigh. dissolves

17th T 105.4 Skin little if at all, injected. Larynx not affected. Inguinal glands enlarged and also one or two, a inch post. to the navel.

Animal looks tolerably well

18th T 102.3 Not nearly so much affected as the other than a little by hyperaemia. no papules. Inguinal

glands swollen and hard.

E T 103. Glands swollen as before, & me on loins
not much skin eruption.

19th M. T 101.3. Caudal about same. E T 104

20th M. T 103°. Very little redness. P M 104

21st. a.m. 104 / P.M. 103.4


22nd. a.m. 102. does not appear very ill


23d 103.5°


24th 103°


25th 104°


26th 105°. does not appear so well has slight purpura
of annus. 27th 104°. 28th 104.









































of 1871
of 1872
of 1873
of 1874
of 1875
of 1876
of 1877
of 1878
of 1879
of 1880
of 1881
of 1882
of 1883
of 1884
of 1885
of 1886
of 1887
of 1888
of 1889
of 1890
of 1891
of 1892
of 1893
of 1894
of 1895
of 1896
of 1897
of 1898
of 1899
of 1900

2000000000

Handwritten text, possibly a name or title, written in cursive.

Handwritten text, possibly a name or title, written in cursive.

Handwritten text, possibly a name or title, written in cursive.

Handwritten text, possibly a name or title, written in cursive.

Handwritten text, possibly a name or title, written in cursive.

Handwritten text, possibly a name or title, written in cursive.

Handwritten text, possibly a name or title, written in cursive.

Handwritten text, possibly a name or title, written in cursive.

Handwritten text, possibly a name or title, written in cursive.

Handwritten text, possibly a name or title, written in cursive.

Handwritten text, possibly a name or title, written in cursive.

Handwritten text, possibly a name or title, written in cursive.

Distinction & et al. per dicitur

Obsequy m. the deduction

Quadrifidum & tunc in

Ritzy 1. red. munda

Respiratione hactenus ecclesiastica

Misogynia & munda

Quintus ponderabilis

Expositione & apponit

Quintus due profer

Quintus apud profer. ^{to what}

Quintus. munda & abund

Quintus tunc. munda. munda

Quintus. munda & munda

Quintus. munda. munda

Wagner's *Wagner*, to include each - transfer

Wagner and down and
Wagner to stick clear

Wagner to refer to, Wagner, Wagner

Wagner to W. Wagner, Wagner
Wagner, Wagner, Wagner (inf. to end)

Wagner to Wagner

Wagner to Wagner, Wagner, Wagner

Wagner to Wagner, Wagner

Wagner to Wagner, Wagner, Wagner

Wagner to Wagner

Wagner to Wagner, Wagner, Wagner

Wagner to Wagner

Wagner to Wagner, Wagner, Wagner

Wagner to Wagner, Wagner

Wagner to Wagner, Wagner

Wagner to Wagner, Wagner, Wagner

Wagner to Wagner, Wagner

Wagner to Wagner, Wagner, Wagner

Experiments forming independent proceedings
conducted to test trials

Experiments on the formation of
liquor of a magnetic glass

Experiments on the formation of
liquor of a magnetic glass

Experiments on the formation of
liquor of a magnetic glass

Experiments on the formation of
liquor of a magnetic glass

Experiments on the formation of
liquor of a magnetic glass

Experiments on the formation of
liquor of a magnetic glass

Experiments on the formation of
liquor of a magnetic glass

Experiments on the formation of
liquor of a magnetic glass

Experiments on the formation of
liquor of a magnetic glass


Handwritten text at the bottom of the page, including the word "Handwritten" and "to provide to meet with".

Handwritten text in the lower middle section, including "Handwritten", "to meet with", and "to provide".

Handwritten text in the middle section, including "Handwritten", "to meet with", and "to provide".

Handwritten text in the upper middle section, including "Handwritten", "to meet with", and "to provide".

Handwritten text at the top of the page, including "Handwritten", "to meet with", and "to provide".



Wahrscheinlich dass

Es ist die - beginnt er mit der

Es ist die Beispiele

Es ist die, welche

George Washington

Every day name

Washington from interest

Washington & the change

Washington, nearly

Washington, spring

Washington land, much with ground

Washington, stable (1000)

Washington, spread, much with

Washington, square below

Washington, thick, very, slope, level

Washington, land, very, man

Washington, some times

Washington, about, with, ground

Washington, get, in, the, building

Washington, work, mine

Washington, in, work

Washington, the, work, & on, the

Washington, & mistake

Washington, considerable

Washington, & carry, out, work

Washington, even, across, the, river

Washington, in, the

Washington, & health

Washington, & interest, in, the

transformation the banking contact
reach of the bank case
efficiency along
Kahn trade lead
Kaufmann's & company
Kaufmann & Co. reach
Kahn's & the direction, arrangement
Kaufmann & arrive at
Kaufmann alle still
Kaufmann's influence
Kaufmann's & change count
Kaufmann & income, expense
Kaufmann & produce, general
Kaufmann's & condition quality
Kaufmann & state the up
Kaufmann's & circumstances condition
Kaufmann's & purchase
Kaufmann & case

transformation the banking contact
reach of the bank case
efficiency along
Kahn trade lead
Kaufmann's & company
Kaufmann & Co. reach
Kahn's & the direction, arrangement
Kaufmann & arrive at
Kaufmann alle still
Kaufmann's influence
Kaufmann's & change count
Kaufmann & income, expense
Kaufmann & produce, general
Kaufmann's & condition quality
Kaufmann & state the up
Kaufmann's & circumstances condition
Kaufmann's & purchase
Kaufmann & case

Steckbrief. Spuk. Spuk.

Sticht f. prof. day. auford.

Stechen f. lat. f. antine

Sticht f. call. (p. 111)

Sticht f. begin. in the first

Sticht. ohne f. auf.

Sticht f. kund.

Sticht. ohne

Sticht f. lat. n. by kein auf - (sammt)

Sticht f. Sticht. auf

Sticht f. same up. left up.

Sticht (n) kann. haben

Sticht. f. auf

Sticht. f. in

Sticht. ohne. Sticht. auf. auf.

Sticht. f. auf. auf. auf.

Sticht. ohne. Sticht. auf.

Sticht. auf. Sticht. auf.

Sticht. (n) Sticht. auf.

Sticht. f. kein. auf. auf.

Sticht. Sticht.

Sticht. es. Sticht. auf.

Sticht. f. Sticht. auf.

Sticht. f. Sticht. auf.

Ant - light blue compound. - on shore by

August 29, addition, several

Went to see the fishery

Program for the

Indians to separate

Worming, original furniture

Wing riding, mowing, in Wagon after

Stay in, mowed, foundation

Quail (in) quail left

Schulze, extra line

Reder (in) have, symbol

Smacken or turning, adj. full grown adult

Engelsen, + diagram which, dried

Scham + share - paid, symbol

Live in base. - choice of material

Moberg's concerns

Moberg's + over + layers

order + m. m. st. (Wagon) drawers

Schickling's shipping, above

Order from separate, to write

Shower of water, glass, ornament

Schweinfurth, in, water, quantity

system in the small

Preservation of - contraction

of - separate, quantity

fast abend. nearly
 Abingen & again. steam. rack
 Festigkeit former
 Rechtigkeit. drossel-composition
 Wichtig important
 Einhalten wrapped up
 Bauen v. t. leiten, t. 4. (over m.) (m.)
 Anfahren & weg. lay hold of
 Anhalten & small up
 Einhalten & drosseln. rarr
 While the case will
 Anhalten & halten
 Einhalten & die Handlen
 Einhalten & halten.
 Anordnenung & t. t.
 Einhalten. (all the case here, also, with)
 Abnehmen & drosseln drosseln
 Wache & case rarr
 Anhalten & rarr.
 Key on (Anhalten) & drosseln
 Einhalten. & rarr.
 Einhalten (op) & rarr
 Einhalten. in der
 Einhalten & rarr. (also rarr. rarr)

Perlygen + hant + rygum

Bemrasschar + cause

Empfindlich ^(Empfinden + feel) sensible ^(Empfindliche)

Bemerkung, dany, performance

Benflache f. surface, superficies

Abtoren fl. covered with + green house, fl. of

Rand on. edge, bank, border

Amphorn + cause + des substance

Breitm + apart, apply

Bruder, less small (Töchter, wachser)

Wäher gum, Wäherg rauen, Plümen

Wannthelbar immediate direct

gerthel- delivery

im Grunde in truth

Amphoren + arranged, advice

Empfänger answer + certificate

Tem. glue, lime

gerben + tan

Schale f. shell, Schalehaut + clamm

linierfal in cartilage graft

Erdeungum + such by gum + puncture

schleimig simile

Wing measuring in Wörtern aufstehend
Wing + inner system
Wing + inner system

Grade 1. Schumann's. part.

Harmon & cell

Trimmer & separate

Large narrow clear top.

Raum in space room

Wimmeltten. stark unimmediate

Gerbrüder. something. main

Entzogen & endow. & abstein from

Shoag (my) end rope string

Kadlmann's. (2) shared child

Wepallik. small red pin before. influence, being

Quarmer. to make and. to sell. double

Unberathung. in dinner

Wimmeltten. unimmediate

Thick party

Zerschell. (2) Professor

Remains. (and. digestive canal

Reife of new seeds like ones

near f. living

Fast almost nearly

Schuldlos. Praxys. (2nd part top)

Sanctus. separate

Wimmeltten. at bottom

Praxys am. Stamm

Andenken. & unimmediate. unimmediate

Stimulans - Charming, the - wgs. d. wgs.

Watrachung / einmischen, enthalten

Aufsatz f. parthen

Ammonien teure parthen

gefallen fast in parthen

Entsch. entke parthen

Reinigt. trank, brennt

Hammerbeile. parthen-paar

schadet, stult

Bauch m. belly, ernst

gelent. n. i. parthen-paar, parthen-wgs

istlich nicht parthenbeile

Ammonien & tier, stornel

gleichmässig, unklar, spalten

Ammonien & tier, stornel

Schmelz. die tier

gabe f. wgs. - wgs. wgs

Wollwurm, parthen, entke

In der that. in d. wgs.

Chinik. die parthen

Empfind. n. unklar

geringfügig, unklar

Mücken parthen, parthen, parthen

gibt lang, tauchen

Antwörung W. a. a. a.

Bestand: partei a. a. a.

Reise, a. a. a. a. a.

Verfahren & a. a. a.

Angewandte a. a. a.

Gründung a. a. a.

Einfluss & a. a. a.

Einfluss (a) a. a. a.

Einfluss & a. a. a.

27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100

Bedienung (m) personal

Bedienung teatry
Befahrung & mack

Bedienung mitfranke

Bedienung theorie kind. theorie. art
Bedienung & provide. (tomm an error
Bedienung & mack. mack. mack. mack.

Bedienung

Bedienung (m) mack
Bedienung (m) mack

Bedienung mack, mack, mack

Bedienung mack

Bedienung mack mack
Bedienung mack mack mack

Bedienung mack (m) mack

Bedienung mack mack mack

Bedienung mack mack

Bedienung mack mack

Bedienung mack mack mack

Bedienung mack mack mack

Bedienung mack mack

Bedienung mack mack mack

Bedienung mack mack

Bedienung mack mack mack

Bedienung mack

Strecken + mit Speise

Frederick in praxi

Waldmann Carbon acid Carbonate

Wilde + Speise Speise

Waldmann + Speise

