

Vol 1

297

Pl 41

Carulocephalus.

25. Ps. Brachyurus rostro nigro angulo

superiore rubro Capite collo et superiore

Nat. p 103

parte pectoris cœruleo sed maculo nigro

infra auribus Ventre Abdomine cauda

et aliis viribus, sed in superioribus partibus

alarum flavescente, ano phenicio cauda

subtus cœruleis. Habitat apud Vera Crux in India
Occidentale. L. S. N. p 100

The Blue Headed Parrot.

111
dum impinguem dicitur
miserabiliter super adiutoria
miserabiliter super adiutoria
miserabiliter super adiutoria
miserabiliter super adiutoria
miserabiliter super adiutoria

310

Vol 1

298

or 42

Melano Caphalus

33 P: Brachyurus Viridis Pileonigro

pectore Albo. L. S. N. p. 102

Habitat in Mexico

103
Nat. p. 103

The Black Headed Parrot

with a White Breast.

Vol 1 ~~300~~ N^o 44

23. *Domicella*

Ptillicus brachyurus coccineis

orientalis vertice nigro rostro

et Iridibus flavis. cera grisea pedibus

albis. humeris ventre et ab domine

caerulecentibus. Cauda exteriore

parte caerulea subtus verde genibus

collo pectore lateribus dorso et

Uropigio coccineis alis supra

viridibus infra coccineis remigibus


supra viridibus infra nigris

macculo flavo notatis. L.S.N.^o p. 100

Habitat in Asia

The Red and Blue Lory

the painted Lory.



310

301


Iris.

Pj. Br. coccineus, alio viridibus
maculis flavis, pectore rostrum flavum,
occipite cyaneo.

Zostrum flavum; Cera nigricans. Caput
sapra antice nigricans prospice
cyaneum. Circulus oculorum nigricans,
Iris tutea. Collum coccineum. Pectus ^{ad Nat. p. 103}
flavum. Dorsum coccineum. Ala
virides: maculis ^{Aumaris complex} flavis; cauda corpore
brevior, truncata, coccinea, apice viridi.
Pede, nigricantes. Temora carnea.

Habitat in India orientali.

The painted Lory.



310

308

302

Pullarius.

Psittacus Brachyurus parvus

Viridis, cauda coccinea fascia

Nigra sed apice viride. Dorso et
Uropigio Cæruleis Splendentibus
in maribus, fronte et gulo Coccineis
in Feminis autem flavis vel
aurantiis.

Habitat in Africa

The guinea Sparrow.

ne

orig.


ad Nat. p. 103

em

tis

si

1



310

308

309

303

Lory Capite Cæruleo
Psittacus Brachyurus Prostro
 flavorubes cente Pilo cæruleo, vertice,
 et genibus coccineis. Iridibus flavis
 a quibus linea nigra protracta est
 Pectore coccineo lineis viridibus
 Transversalibus ventre viride. Cauda,
 et ab domine flavis. alis et superiori
 parte viridibus altero sexu viribus
 flavo variegato.

The Blue Headed Lory

orig.
ad Nat. p. 103
em

John D. King, M.D.,
Montgomery, N.Y.
July 11, 1862.

310

308

309

Lory Coccineo
Capite Nigro.

309

304


Psitacus Brachyurus puto nigro
Corpore coccineo. alis viribus. Rostro
alto.

wings

ad Nat. p. 103

The Scarlet Black headed,

Lory



310

308


304

305

Lory Coccineus

Psittacus Brachyurus rostroflavo
et Nigro, Capite ventre et Uropigio
Coccineis Humeris flavis. Dorsso
purpureo maculo flavo in medio
posito. alis et femoribus viridibus;
pedibus Nigris.

The Purple Lory



310

308

309

306

7

Pitticus Capite Cœruleo
Pectore fusco

or

Pitticus Brachyurus rostro cinereo

2d


Capite Cœruleo Dorsò Viride Remigibus
Cœruliis Pectore pallide fusco femoribus ^{moris}
Luteis

ad Nat. p. 103
nem

Habitat in Hispaniola

is
tri

The Brown breasted
Blue Headed Parrot



310

Pittacus cyanocephalus.

309
7

307

Galgulus.

or

2d

36 Pittacus Brachyurus Viridis

Uropigio pectorq; in medio.

ee

coccineis. vertice cærulic. Linn. Syst. N.

meris.


Habitat in India p. 103. 36. nem

ad Nat. p. 103

The small Blue headed

ii,

Parrakeet with a red spot on the Breast.



Pittacus cyanocephalus.


309

307

P. brachyurus viridis, capite coruleo, cauda
fusca basi rubra, rostro nigro angulo baseo
rubro.

Rostrum nigrum: Mandibula superiore ad Nat.p.103
angulum inferiorem baseo rubra. Caput
Collumque corulea: macula nigra utrinque
pone aures. Corsae viride. Alæ virides humeris
flavescensibus: Remiges primariae superne ad
marginem interiorem & inferne ad marginem
exteriorum nigricantes. Cauda aequalis:
Rectrices 12, superne lateribus coruleis,
intermedio viridis, subbasim omnibus basi
rubris apice viridis. Pedes lividi.

Habitat in Insulis Granadis & in
Vera Cruce India occidentalis.



310

Pithecus cyanurus.

309

309

Pitacus Benghalensis minimus
Pitacus Brachyurus minimus
veridis vertice flavo rostro nigro coccineo

Habitat in India Orientale in Bengala

The Smallest yellow headed
Parakeet

Nat. p 103

100

John C. Frémont
and his party
in California.


Pittacus cyanourus.

*P. brachyurus viridis, capite, collo, maculaque
alarum flavis, dorso nigricante, cauda
rubro violacea.*

Magnitudo *P. pullarii* Linii. *P. Loxia curvirostra*
Caput & Collum e viridi flavum. Rostrum
pallidum: cera fuscata. Nares alle in cera
porosa. Pectus & Abdomen pallide viridia.
Dorsum nigrum, interdum paulo virescens.
Ale supra nigricantes macula magna
obliqua flava: marginibus corulescentibus.
Pes: Margines remigium primorum
 virescunt. Ale subtus corulescent. Cauda
angularis, e rubro violacea: Remigibus
 acuminatis lateralibus prope apicem
 macula nigra notatis: intermediius duabus
 immaculatis. Pedes pallide fuscui. Pes: Tectrices
Caudae subtus virides. Angulus inferior
humerorum miniatus.

Habitat in Insulis Granadis.

Nat. p 103



310

309

Pittacus granadensis.

P. brachyurus viridis capite nigro, collo superne
auransiaco, cauda subtus basi flavecente.


Caput totum nigrum. Prosternum albicans.

Collum superne ferruginea auransiaca,
inferne viride, subtus viridi lutescens.

Corpus viride. Ale virides. Cauda aequalis,
viridis, apice corulescens, subtus basi
flavescens. Pedes pallidi.

Nat. p 103

Habitat in Insulis Granadis.



310

Ramphastos

*Piostrum. maximum, inane, convexum.
extorsum serratum,*

Nares. pone maxillas

Lingua pennacea

Pedum digiti anticiposticq; gemini. Lin. Sys. Nat. p 103

The Toucan

U. M. D. P. M. D.

Parvulus, pectus, conatum, mordet
mollis, mordet
pedem, mordet,

mordet, impul

pedem, mordet, impul

mordet, mordet


310

Ramphastos nigrirostris

P: rostro nigro: carina compressa, gula lutea
fascia pectoris eropugio anaque sanguinea.

Prosternum nigrum, basi album, carina compressa:
Mandibula superiore in aliis ferrata in
plerisque integra. Caput & Collum supra,
¹⁰³ Dorsum, Abdomen, Alae & Cauda nigra. Gula
ad Basin mandibulae inferioris lateribus alba,
in medio aurantii coloris usi etiam totum
collum sublatus. Fascia in pectore sanguinea.
Anus & Uropygium sanguinea. Penniger
20. Rectrices 10.

Habitat in Insulis Granadis.



Ramphastos

Piperivorus.

*Ramphastos rostre nigro carina
erassissima. L.S.N. p. 103*

Habitat in America meridionali.

The Black Beaked Toucan.

from the Island of the Granades

W. H. Dugay, Esq.,

Montgomery,

Mass., May 10th, 1842.

Dear Sir, I enclose you

sums of \$1000, \$1000, \$1000,

which you will find correct.

Ramphastos

Tucanus.

Ramphastos rostro rubro: carina obtuso albo.

*Habitat in America Meridionali
Maxilla versus basin fascia nigra*

The Red Beaked Toucan
from the Island of γ^e Granades

Montgomery

Montgomery

After much musing when others had disagreed

Montgomery agreed with all
and made much noise about it.

Montgomery, April 2d,

Montgomery, April 2d,

314

312

Ramphastos Tucanus.


R. rostro rubro: carina obtusa alba, corpore
nigro gula flavescente fascia anogit san-
guinea, uropygios flavo.

Linn: Syst. Naturae. p. 103. 2.

Caput & Collum supra nigra ut & Dorsum-
Ale, Cauda & Abdomen. Gula & Collum sublata
exalbido flavescentia, fascia in pectore san-
guinea. Anus croco-sanguineus. Uropygium ut
flavum. Pedibus laterales breviores.

Habitat in Insulis Granadis.

n
ed
er



314

Ophryosaurus Aracari.


*A. rostro nigro: maxilla superiore lateribus alba,
abdomine sulphureo fascia uropygialis sanguinea.*

Linn: sp. n. p. 104. n. 4.

*Synonymum Edwardi hoc non pertinet.
Caput, Collum, Dorsum, Alae & Cauda nigricans,
fascia & paulo e viridi nitent Abdomen e viridi re-
glavum, in medio fascia sanguinea. Uro-
pygium sanguineum. Anus colore abdominis,
taenio autem uropygii longiores interdum la-
teribus dependentes subrubescentem reddunt
Anum. Prostria mandibula superior basi
triloba, lateribus alba, canina nigra obtusissi-
ma, interdum serrata plerunque levis.
Mandibula inferior nigra*

Habitat in Insulis Granadis.

*soon
when he returned
to the trial of his time keeper
covery of the Longitude.*



Pramphastos

Aracari.

*Pramphastos rostro nigro: maxilla superiore
lateribus alba; basi triloba.*

Toucan Edw. av. 64 t. 64

aracari/margrave Cras. 217 Rai. av. 44

Habitat in America meridionali.

maxilla superior ad nares triloba,
quod non in reliquis. fascia abdominis
anoq sanguinis. L S N. p. 104

The Toucan

with a Black Beak striped with White.

N.B. These Birds came from the Islands called the
granades. & were brought me by M^r W^m Harrison
with many others in y^e year 1764 when he returned
from Barbadoes after the trial of his time keeper
for the discovery of the Longitude.

estudium.

mentis.

propositum. Atque in omnibus istis studiis
ad hanc modum. Non obstat
propter ipsas. sed propter omnes
dilectionem fratrum et similitudinem.
aliquot annos haec res ipsa
conducit ad. Similiter et res huius
propter. hanc similitudinem. pone

mentis. sibi.

Atque. Miserum. Quod deinde a filio
ad hunc etiam. ut non esse debet. Atque
miserum. quod per se deponere. est. Atque
deinde. cum. pater. non. potest. non. alii
magis. cum. filius. deponere. non. potest. non.

315

Ramphastos

Picatus

Ramphastos. rostro rubro: apice nigro
carina compressa.

Picus Brasiliensis: Alb. av. 2 p 24. t 25 Aldr. orn. 620.
c 19.

Habitat in America meridionali

W. M. Thompson

Editor

newspaper editor and publisher

Montgomery, N.Y.

and author of "The Slaveholder's List."

"An American Slave and his Master."

315

AVES PICÆ RHAMPHASTOS

S. MOMOTA

Rhamphastos pedibus gressorius,
rectricibus duabus intimis medio denuda-
tis L. S. N. Ed. 12. p. 152.

MOMOT Will. orn. 295. Raj. av. Ed. n. av. t. 328
Briss. av. 4. p. 465. t. 35. f. 3.

Gura Guainumbi. Margr. bras 193. Rei av. 49.

СОТЫИ МАНГАДЫ

АГАМЫ

317

315

Aves Picae


Buceros.

Rostrum convexum, cultratum magnum

Frontis calvaria nuda, oeso-gibbosa.

Nares. in Basi rostri.

Lingua acuta, brevis.



Buceros 315

2. Rhinoceros.


Rhinoceros. Buceros cornu maxillari
frontis Recurvato.

Habitat in India.

The Rhinoceros bird.

This bill is the 2^d sort in Willoughby's ora.
Table 3. it varies from Willowby's descrip.
in size. this Bird being abt of magnitude
of the Raven. in Laving the Whole body
black except of Highs the Ends of ye
tail feathers & feathers under the tail
only White.

The Legs are covered with short black
feathers down to the feet.




57

317

316

Crotophaga

Rostrum Compressum semi ovatum
sulco utrinq; exaratum; Mandibula
superiore marginē utrinque angulata
Nares per rectas.



57

317

316


Ani

*Crotophagus ater Rostro breviore
compresso acuato cultrato*

Linn. S. N. p. 105. 1.

Habitat in America

The Brasilian Ani




317

Torquilla sive Ipx.

Cuculus subgriseus maculatus, rectricibus
nigris fasciis undulatis. L.S.N. p. 112.

The Wry Neck



320

318

Synax granadensis.

I. nigricans fronte gulaque coccinea, *abdomine* flavescente.


I. Torquilla paulo major. *Rostrum* nigricans rectum: *Mandibula* superiore ultrata, tereti.

Caput nigricans, anteriorius paulo virescens: fronte coccinea; linea albida supra oculos. *Collum*, *Dorsum*, *Ale* & *Cauda* nigricantia, plumis viridiibus hinc inde interspersis, praecipue in aliis ubi linea obliqua flavicans. *Gula* coccinea. *Pectus* & *Abdomen* flavicantia. *Pectrices* apice obtusa. *Pedes* nigri: digitis anticus duobus prosticisque duobus.

Oc.: Prope narum seta nonnulla nigra. In mandibula inferiore fascia videtur obliqua, pallida.

Habitat in Ovulis Granadis.

The facanacori of Margraf Willoughby p. 139.
seems to be of the same genus.



320

319

Auratus.

*C. cauda subforcipata, gula petore que
nigra, rumpa rubra, racis remigum*

primorum aurea. L. S. N. p. 112
Habitat in Carolina.

The golden Woodpecker.

lathes
impaling him. After which, when I
had no more to do, I went down
to the village, and there I saw
the old man, who was still
standing by the roadside.

II 320

320

Pileatus.

P. niger, capite cristato rubro, temporibus

alisque albis maculis. L. S. N. p. 113

Habitat in America

Amulius.

Amulius, whom deducit deus impud.

Amulius.

Amulius, whom deducit
deus impud.


320

Martius.

Picus niger, pileo coccineo. L.S.N.p 112

Habitat in Europa.

The Black Woodpecker



320
321 (VJD)

Principalis.

P. niger, crista coccinea, linea utrinque
collari remigibusque secundarisi albis.

L. S. N. 113

The largest Black Woodpecker
of Carolina

Galapagos

Galapagos Islands

Galapagos Islands

Galapagos Islands

322

Picus Pileatus

*Capite Cristato Lutea,
temporibus Luteis*

*Habitat in Carolina
Famina precedentis ut opinor*

The Yellow Crested Black
Woodpecker

Family, family,

Mother, Mother, mind
Mother, mother, mind

Mother, mother, mind.

Mother, mother, mind.

Mother, mother, mind.


D

326

373

*Picus Viridis Capite
Cinereo.* Edn^o p. 65

Habitat in Norvegia
The Grey-Headed Green
Wood Pecker



D

326


324

Europaea.

*Sitta. rectricibus nigris: lateralibus quatuor
infra apicem albis.* L.S. N.p. 115

Habitat in Europa.

The Nuthatch.



D

326


328

Epopos.

1. *Upupa. cristata variegata.* L.S.N.p. 117

Habitat in Europæ Sylvis.

The Horope.



D

326

326
Viridis.

P. viridis, vertice coccineo. L.S.N.p 113

Habitat in Europa

The green Woodpecker

77

R
Silius.
Silius. min. mod. 1.
Silius. min. mod. 2.
Silius. min. mod. 3.

326

Benghalensis

*Picus viridis, occipite rubro, nucha
nigra, subtus anticeque albus nigro
maculatus.*

L.S.N.p. 113

The Crested Green Woodpecker r

inventus puer.

ab aliis sedem alii per latitudinem ambi.

vixit etiam impeditus videtur. Vixit

etiam.

. latitudinem

inventus puerus videtur.

327

Major

*Palbo nigroque varius, ano occipite
qrubro.* L.S. N^o. 114

Habitat in Europa.

The Great Spotted Woodpecker

Digitized by srujanika@gmail.com

328

Minor

P. albo nigroque varius vertice rubro,

ano albido.

L.S.N. p. 114

Habitat in Europa.

The Least Spotted Woodpecker

W.M.

329

Tridactylus

*P. albo nigroque varius, pedibus
tridactylis.* L.S. N.p. 114

Habitat in Siccia

The Three Toed Woodpecker.

John Smith
writing upon paper
in his library
and in his
writing book.

Picus ferrugineus

P. cristatus ferruginosus, cauda nigricans.


Magnitudo P. medii Linn: Totus rufescens ferrugineus, sed Caput supra & Pennae longiores
qua cristam formant, parum lutescent.

Tectrices alarum maculis paucis pallidis
adspersae. Alio sublissus fusco fasciis dilute
ferrugineis. Urofygium lutescens. Cauda
nigricans acuta: Rectricibus decem,
acuminatis, intermedio longioribus.

Prosternum flavicans. Pedes livido-fusca.

Habitat in fructu Granadis.

The Chestnut colour'd Woodpecker)



Picus granadensis

*P. e fuso viridis, occipite nigricante, nucha late
ribusque gulo coccineis, abdomine lutescente
fasciis transversalibus.*

" *Pico minori Linn: paulo major. Caput supra
e plumbos-nigricans, frontice coccineum; Regio
oculorum albida, fruis fuscis; Gula latera prope
rostrum coccinea. Collum supra, Dorsum, Ala
& Cauda e fuso viridia seu subcuprea. Collum
sublatus, Abdomen & Pectus lutescentia fasciis
numerosis transversis fasciis. Cauda apice
nigricans. Postuum & Pedes nigricantes.*

Habitat in Proculis Granadis.

Wendy will

333

P
Pyrrhocorax

U. atra, rostro pedibusque rubris. L.S.N.p. 110

Habitat in Anglia,

The Cornish Chough.

commodius

modus

modus

modus

Egyptica.

U. nigra Cœrulecente major, ^{in dibus floris} dissertation
 a precedente in magnitudine. et rostro; quid
 est maius. arcuato. et Naribus longioribus
 nudis.

Habitat in Agypto
 The Egyptian Chough.

188

W. H. C. -

W. H. C. -

W. H. C. -

W. H. C. -

335

Familiaris.

1. Certhia. supra grisea, subtus alba,
remigibus fuscis: decem macula alba. L.S. N^o. 110

Habitat in Europa

The Creeper.

William

and a willow, many aged willows
and alders, all about the river which
is broad in width.


William

Alyanthes 336

Certhia coerulescens dorso superiori, aliis caudae nigrae, vertice
cyaneo azureo.

luisine annatum

Descrip. Rostrum, pallide nigrescens. Caput picea splendidè
azureum, maxilla inter basim rostris & oculos nigra. Tota
avis fimbriae coerulescens, dorsum superius, alæ & cauda nigra.
Ullam remiges intorsum ad marginem lateris. Pedes
lutescentes. Magnitudo Certhia familiaris.



Cruentata.

O. nigro-cærulescens, subtus alba, vertice.

cervice, dorso uropygione rubris. L. S. N. p. 119

Habitat in Bengala

The Red Creeper.

