

Note on Lophoceros from  
Yukon Territory, Dec 1884.

1. Specimens, <sup>mostly</sup> in white siliceous stone  
Lewis River.

*Taxodium Tinctorum*, Heer

*Glyptostrobus Supraeus*, Heer  
They in black shaly  
rock from Runk Rapids

*Sequoia Langsdorffii* Heer  
Some fragments but  
very certain.

The above three Conifers  
are found at English Bay and  
at places near Sitka in  
Alaska. (Fl. Foss. Alaska 1869)

*Equisetum*, species

This is similar to one  
figured from various northern  
determinable, in gray micaceous  
sandstone, Coal locality Lewis  
River Ogilvie's collection.

2. Specimens from Pelly River  
in black carbonaceous  
shale,

The few specimens examined  
are full of impressions of *dictyodendroides*  
coniferous leaves. Much  
crushed and imperfect,  
one has the venation of  
*Corylus ellae Guerinii*, Forbes  
Another seems to be *Inglauis*  
near to *I. acuminata*, Braun  
Both of these species are  
said to have been seen at  
English Bay Alaska, and  
also at the Kenai River.  
The determinations cannot however  
be considered certain.

3. Fruit Woods, Park Rapids,  
Lewis River,

There are two species both well  
preserved. One may be a specimen  
of the *S. gigantea* type. The  
other is apparently a *taxodium*  
tree, possibly a *Salisburya*.

All of the above fruits are referable  
to the *Fertig* (Svensen & Jarman) group;  
and if *Cyatocarpus* must belong to the  
extreme upper part of that system.

Merino Plants

Leino

Polly R

Good Cal

Dec 1887