

GEOLOGICAL NOTES.


[The bearings refer to the magnetic meridian.]

- 61-61. The section across the Huronian rocks on the St. Lawrence, shows a variety of dark grey, hornblende and epidioritic schists, dark green massive and schistose diorites, argillaceous reddish diorites, grey and greenish fine grained quartzose mica-schists, and fine grained hard grey imperfect gneisses. The general strike is south-westward.
- 62. Dark green diorite and dark-conglomerate, with a hill to the northeast consisting partly of grey and partly of red fine grained granite.
- 63-63. The Huronian rocks exposed along the Black River consist of dark greenish green and reddish diorites (some of them porphyritic), red and grey granites, conglomerates, fine grained mica-schists, diorites, siliceous, argillaceous, hornblende, chloritic, felspathic and epidioritic schists and imperfect gneisses, similar to those on the St. Lawrence. The general strike is W. S. W.
- 64. Greyish and reddish gneisses, occasionally interstratified with bands of dark hornblende schist and very light grey gneiss.
- 65. Fine grained dark green hornblende schists, having the same strike as the adjacent gneiss.
- 66-66. Hornblende, siliceous and chloritic schists. Strike, N. N. W.
- 67. Huronian schists with shales of gneiss and trap, and veins of calciferous and quartz containing yellow and siliceous sulphates of copper, iron pyrites, blende, galena and malachite.
- 68. Michipicoten Island and the extremity of Cape Gargantua consist of amygdaloidal, conglomerates, sandstones, calcareous limestones, etc., resting on the Keweenaw (Cambrian) series.
- 69. Light red granite of medium texture. Dip from the east by hornblende schist.
- 70. Soft green slate holding layers and patches of calciferous and gabbro of reddish granite.
- 71. Greyish slaty diorite with thick beds of reddish siliceous rock; also impure hematite interstratified with grey siliceous layers.
- 72. Dark coloured hornblende mica-schist.
- 73. Massive green dioritic schist.
- 74. Massive and slaty diorites and grey calciferous schists with traces of copper.
- 75. Massive red and grey granite of medium texture and good workable character.
- 76. Cliff 500 feet high of green slaty diorite with radiating strings of calciferous and quartz.
- 77. Glossy green mica and hornblende schists, quartzite-grey slaty quartzite, massive crystalline green diorite, dioritic schist, soft calciferous grey mica-schist and brittle reddish yellow siliceous or cherty schists. At the N. W. extremity of Cap Chouart a low patch of thin-bedded red and grey sandstone rests unconformably upon these rocks.
- 78. Small patches of red slaty volcanic breccia resting upon gneiss.
- 79. Massive, even and rather finely grained pinkish-grey granite.
- 80. Green chloritic, grey felsitic and fine grained mica-schists, with diorites and gneisses.
- 81. Red and grey syenitic granites.
- 82. Chloritic and dioritic schists with masses of fine-colored granite and syenite.
- 83. Fine grained massive soft greenish-grey quartzite.
- 84. Green hornblende schist with pebbles, chloritic and dioritic schists.
- 85. Narrow bands classed as Huronian, and consisting of mica-schists, hornblende, chloritic, siliceous and felsitic schists.
- 86. Dark grey siliceous slate and dark green finely crystalline hornblende-schist, with greyish mica-schist towards the gneiss on either side.
- 87. Siliceous felsitic and mica-schists, with quartzite and diorites.
- 88. Dark hornblende and mica-schists with layers of light-colored granite running with the strike.
- 89. Beds of lignite, from 1 to 6 feet in thickness in boiler clay. (See p. 4c. Report for 1877-78.)
- 90. Gypsum in the N. W. bank begins at 8 miles, and is not seen in the S. E. bank at 16 miles below this point. It is about 10 feet thick, and is associated with earthy greyish and buff-colored magnesian limestone and calcareous shale of Devonian age.
- 91. Siliceous hornblende schists.
- 92. Red and reddish grey syenitic granite with patches of grey amygdaloid, containing calciferous, fluorapatite and apatite.
- 93. Rather dark-colored hornblende schists with a general westward strike.
- 94. Lead and copper ores discovered by Mr. Brown in veins in Huronian rocks.
- 95. Huronian rocks striking north of east, probably near their junction with the Laurentian.

Geological Survey of Canada.
Alfred C. Sibly, LL.D., F.R.S., Director.

MAP SHOWING APPROXIMATELY THE GEOLOGY OF THE BASIN OF MOOSE RIVER AND ADJACENT COUNTRY. TO ILLUSTRATE DR. BELL'S REPORTS OF 1875, 1877, 1881.

Scale 6 Miles to 1 Inch
Drawn by L. N. RICHARD, B. A. Sc.
1883.
SCALE OF COLOURS
LAURENTIAN. GRANITE. HURONIAN.



GEOLOGICAL NOTES.

[The Bearings refer to the Magnetic Meridian.]

- 1. Felsite, generally ribboned purple and green; some parts grey and siliceous. The general strike is parallel to the course of the East Branch of the Moose River. At the outlet of White Beaver Lake it is associated with a beautiful siliceous brecciated chert.
- 2. Quartzites, mostly light grey, and in places approaching the character of sandstones, which in others they are full of pebbles. Some portions are brownish, some greenish, and others dark grey. At White Beaver Lake the surfaces of the beds occasionally show ripple-marks.
- 3. Massive crystalline greenish grey diorite, mostly of dark shades.
- 4. Dark greenish grey argillaceous mica-conglomerates; pebbles mostly of red syenitic granite, with some of white quartz and green schist.
- 5. Dark hornblende schists, full of veins and masses of grey pyroxene; all cut by veins of siliceous quartz.
- 6. Great alternating masses of grey quartzite and dark-colored diorite, both of which are frequently of a coarse conglomerate character, the pebbles consisting of syenitic granite, quartzite, and various slates of the Huronian series, mostly rounded and crowded together in great basins. These massive beds are interstratified with sandstones at intervals. The dip is from E. to E. N. E., at an angle of 45°.
- 7. Beds of quartzites with copper pyrites and crystalline specular iron. Dip N. 70° W.
- 8. Dark and grey syenitic granite, mostly coarse in texture.
- 9. Dark hornblende and siliceous schists.
- 10. Greenish grey dioritic mica-conglomerate with smooth rounded pebbles of syenitic granite.
- 11. Banded reddish quartzite, mostly fine grained and interstratified with greenish siliceous slates.
- 12. Greenish and greyish argillaceous and siliceous slates.
- 13. Dark green serpentine with crystals and calciferous, and containing chromite ore.
- 14. Massive light green felsitic volcanic ash with dark and light spots.
- 15. Soft greenish grey magnesian rock, rarely stratified and thickly intersected in all directions by calciferous strings.
- 16. Hornblende and dioritic schists dipping south-westward at high angles.
- 17. Greenish grey quartzite, partly massive and partly schistose. They hold transparent grains of quartz.
- 18. Dikes of crystalline dark greenish grey diorite.
- 19. Dark grey hornblende schist. Strike, S. 75° E.; angle northward, 65°.
- 20. Grey massive mica-schist, chloritic and greenish grey siliceous and dolomitic schists, with rusty surfaces and cut by quartz veins.
- 21. Siliceous grey slates, running W. N. W.
- 22. Fine grained greenish grey quartzite, followed, half a mile to the N., by bluish grey slaty-slate, and further N. by more quartzite.
- 23. Straight greenish grey crystalline somewhat schistose diorite, cut by irregular quartz and calciferous veins. The dip is northward at an angle of 80°.
- 24. Grey siliceous slaty, grey felsitic and dark bluish grey clay-slate. Dip N., angle 60° to 70°.
- 25. Green and grey massive chloritic and talcoid schists. Dip N., angle 60°. Cut by a post-dike running N. and S.
- 26. Massive grey semi-crystalline slates, holding grains of specular iron, and cut by small veins of blue-quartz.
- 27-27. Massive and slaty greenish grey diorite.
- 28. Dark grey quartzite and massive light greenish grey siliceous schist. Strike, N. 75° to 80° W.
- 29. Hard greenish grey slaty diorite, cut by a great dyke, running N. and S.
- 30. Soft greenish grey calciferous talcoid schists. Strike N. 55° W.
- 31. Grey volcanic ash, followed by greenish green chloritic and dioritic schists. Dip E. N. E.
- 32. Dark greenish grey hornblende schist, with some chloritic schist. Strike, N. 55° to N. 70° W.
- 33-33. Laurentian gneiss begins at the foot of Davis' Rapids and is found all along the river to the foot of the Long Portage, which appears to mark the southern boundary of the Devonian area.
- 34. Soft dark nearly slaty.
- 35. Grey and buff fossiliferous limestone.
- 36. Soft yellowish and reddish grey porous limestone.
- 37. Calcareous-siliceous chert-like sandstone, with green spots.
- 38. Dark grey crystalline mica-schist, cut by dikes of dark, compact diorite.
- 39. Greenish calciferous mica-schist, mica and hornblende schist.
- 40-40. Tender grey mica-schist with rusty partings.
- 41. Fine grained greenish grey calciferous diorite.
- 42. Grey slightly siliceous argillaceous schists, with thin partings of calciferous in the vein.
- 43. Unstratified greenish grey diorite with disseminated grains of iron pyrites, and in some places holding vesicles filled with calciferous and white quartz.
- 44. Dark green argillaceous weathering dull white, strongly magnetic and containing grains of chromite iron.
- 45. Mica-schist, hornblende, chloritic and dioritic schists, quartzite and diorites.
- 46. Impure, hard, blue-grained grey limestone.
- 47. Obscurely stratified dark greenish grey dioritic schist.
- 48. Hill 600 feet high of fine-grained grey quartzite.
- 49. Conglomerate with pebbles, and slaty matrix with rounded pebbles of greenish diorite, chert and white quartz.
- 50. Hornblende and mica-schists.
- 51. Mica-schist, chloritic and talcoid schists, associated with massive talc, quartzite, magnetic pyrites, iron ore and mica.
- 52. Very light grey and bluish granites associated with dark syenitic granite holding mica and hornblende.
- 53. Greenish green hornblende schist conformable with the gneiss to the east of it.
- 54. Slaty diorites, with layers of ribboned quartzite, jasper and magnetic iron.
- 55. Bed of grey felsitic mica-schist with copper and iron pyrites just below the 7th portage from Lake Timicameung.
- 56. Soft greenish grey siliceous rock.
- 57. Hard dark grey slaty mica-schist.
- 58. Boulder conglomerate overlain by light grey limestone, all belonging to the Niagara formation. (See GEOLOGY OF CANADA, p. 334.)
- 59. Reddish syenitic granite composed of red orthoclase and white quartz with a small quantity of green hornblende.

This area appears to be occupied by massive crystalline dark from diorites, but the central portion has not yet been explored. See Geological Survey Reports 1878, pages 108, 125 and 128; 1877, pages 302 and 312; and 1877, page 57 C.